

Regulation of Lipid Catabolism

Dave Bridges

June 9, 2026

This unit will discuss how lipids are oxidized and converted into energy. Free fatty acids contain substantially more energy per molecule than either carbohydrates or proteins. For more details on this process see Chapter 27 in *Biochemistry: A Short Course* available in reserve¹.

¹ John L Tymoczko, Jeremy M Berg, and Lubert Stryer. *Biochemistry: A Short Course*. W.H. Freeman and Co, New York, NY, 2015

Contents

<i>Learning Objectives</i>	2
<i>Lipolysis Liberates Fatty Acids from Triglycerides</i>	2
<i>Carbohydrate Overfeeding and Lipid Utilization</i>	3
<i>Fatty Acid Oxidation</i>	3
<i>The Role of CPTI in Lipid Oxidation</i>	4
<i>Lipid Oxidation in the Mitochondria</i>	5
<i>Alternative Fatty Acid Catabolism</i>	5
<i>Determining the Energy Content of a Fatty Acid</i>	6
<i>Ketolysis</i>	7
<i>Regulation of Fatty Acid Oxidation</i>	7
<i>Short-term Regulation</i>	7
<i>Transcriptional Adaptations for Fatty Acid Oxidation</i>	8
<i>Reflection Questions</i>	8

Learning Objectives

- Explain how triglyceride breakdown into glycerol and free fatty acids is controlled in adipocytes by hormonal signals.
- Explain how high carbohydrate diets affect fuel utilization, including effects on lipid fuel utilization. Describe at an endocrine level how this is thought to occur.
- Determine how much energy, in ATP equivalents, is released during the oxidation of a given fatty acid. Be able to relate the energy content of a fatty acid, in general to its physical properties (length and saturation).
- Explain the rate limiting steps of lipid oxidation.
- Explain how ketone bodies are converted to ATP in non-hepatic tissues, and what governs this specificity.
- Demonstrate an understanding of how *de novo* lipogenesis and β -oxidation are reciprocally controlled.
- Describe how very long chain fatty acids are oxidized differently from long chain fatty acids.
- Explain how odd-numbered fatty acids are catabolized, including the importance of vitamin B12 in this process.
- Evaluate the role of transcriptional regulation and long term adaptations to fatty acid oxidative capacity.

Lipolysis Liberates Fatty Acids from Triglycerides

Fatty acids are generally stored as triglycerides, and those triglycerides are primarily stored in adipose tissue. There are two main sources of fatty acids, the dietary fatty acids liberated from chylomicrons by the activity of lipoprotein lipase², and fatty acids liberated from adipose tissue³. This step, the conversion of triglycerides to fatty acids is known as lipolysis.

IN ADIPOCYTES THE MOST POTENT ACTIVATORS OF LIPOLYSIS are catecholamines such as adrenaline and cortisol. Adrenaline functions to rapidly activate adipocyte triglyceride lipolysis to glycerol and fatty acids via the activation of two enzymes, adipocyte triglyceride lipase (ATGL) and hormone-sensitive lipase (HSL). These two enzymes work primarily on triglycerides and diglycerides, respectively⁴. Via PKA-dependent signaling, both HSL and ATGL have

² This will be described in more depth in the lipid transport lecture

³ A quantitatively small amount of lipids may be stored in liver (too much of this is called hepatic steatosis) and muscle (intramuscular triglycerides) tissues as well

⁴ The last step, the conversion of monoacylglycerol to glycerol and a fatty acid, is catalyzed by monoacylglycerol lipase, an enzyme not thought to be regulated by hormones.

increased rates of activity resulting in fatty acid release for use as energy in peripheral tissues. Cortisol also increases lipolysis, potentially through several mechanisms, one of which is increasing the levels of ATGL, in a slower more permanent manner.

Carbohydrate Overfeeding and Lipid Utilization


Insulin potently restricts adipocyte lipolysis. The mechanisms of this are still not clear, but at least part of the actions of insulin are to inactivate the lipases HSL and ATGL by reversing the phosphorylation events caused by adrenaline/PKA. States of insulin resistance such as type 2 diabetes means that insulin is less able to keep lipids in the adipose tissue. Interestingly, studies where *carbohydrates* are given in large quantities as part of overfeeding experiments lead to fat accumulation [Acheson et al., 1988]. This was originally thought to be due to activated *de novo* lipogenesis, but more intricate studies showed that instead, the body was adapting to high levels of carbohydrates by sparing fat [McDevitt et al., 2001] and that the level of lipogenesis was minimal. This lipid sparing effect is the result of increased glucose oxidation and reduced lipolysis and β oxidation. This is a concept we have discussed several times already in this class, when one macromolecule is in relative excess, we tend to alter our metabolic pathways to use this for fuel⁵. Of the multiple potential mechanisms for this lipid sparing effect, one is that carbohydrates induce a strong insulin response, which prevents lipolysis and results in lipids being trapped in adipose tissue. This distinction is key to understanding how high carbohydrate diets relate to obesity.

⁵ Some examples we have already discussed include the inhibitory effects of Alanine on Pyruvate Kinase, Citrate on PFK1 and Palmitate on Acetyl-CoA Carboxylase.

Fatty Acid Oxidation

Fatty acid oxidation is important in many contexts, including during endurance exercise, and fasting. It is particularly important in the heart where even under resting or elevated carbohydrate conditions, the majority of ATP is derived from fatty acid oxidation [Neely and Morgan, 1974]. Once fatty acids enter or are made available to the cell, they are quickly converted to "activated" forms known as acyl-CoA molecules⁶. This is also the first step for triglyceride esterification, and consumes two ATP equivalents. The enzymes that catalyze these reactions are known as Acyl-CoA Synthetases. More details about these enzymes can be found in this recent review: [Grevengoed et al., 2014].

⁶ The acyl refers to a generic fatty acid molecule, so the specific end product could be oleyl-coA (from oleate) or palmitoyl-CoA (from palmitate).




While there are several Acyl-CoA Synthetases, one in particular,

ACSL1⁷ seems particularly important for fatty acid oxidation, as its genetic ablation in mice almost completely prevents fatty acid oxidation [Ellis et al., 2011]. Interestingly, ACSL1 is present on the surface of the mitochondria in a physical complex with CPTI, the most important regulatory protein in fatty acid oxidation [Lee et al., 2011].

MITOCHONDRIA ARE KEY FOR FATTY ACID OXIDATION. Like most amino acid catabolism, fatty acid conversion to energy requires mitochondria. However, unlike both pyruvate and most amino acids, the transport into the mitochondria is the rate-limiting step for fatty acids. This transport⁸ is controlled by the carnitine shuttle system.

The Role of CPTI in Lipid Oxidation

The carnitine shuttle system is a mechanism to get long chain fatty acids⁹ through both the outer and inner mitochondrial membranes. It comprises of two reactions and a transporter. The reactions are overall energetically neutral, with the end products identical to the initial reactants. The two reactions are catalyzed by Carnitine Palmitoyltransferase I (CPTI):



and Carnitine Palmitoyltransferase II (CPTII):



There is a transporter on the inner mitochondrial membrane called Carnitine Acyltransferase (CAT). This transporter *only* works with the carnitine conjugated fatty acids, so this transport system requires carnitine for its activity. As we will discuss in the non-protein compounds made from amino acids unit, carnitine can be made endogenously from the essential amino acid lysine. Carnitine can also be obtained in the diet, with particularly high levels in red meat.

MALONYL-COA INHIBITS CPTI. This is the most important regulatory step in fatty acid oxidation. If Acetyl-CoA Carboxylase (ACC) is activated due to an excess of Acetyl-CoA, Citrate or insulin signaling¹⁰, then there is a buildup of its product Malonyl-CoA in the cytoplasm. This product potently inhibits CPTI [McGarry et al., 1977]. The result of this is to prevent fatty acid import into the mitochondria and therefore fatty acid oxidation. The activated fatty acids are then shuttled towards triglyceride synthesis rather than oxidation. This regulatory process is illustrated in Figure 1. The flip side of this inhibition is that when AMPK or PKA¹¹ are active, ACC is inhibited

⁷ Abbreviated from Acyl-CoA Synthetase Long Chain Family Member 1, indicating a preference for long chain fatty acids

⁸ Which is more difficult than passive diffusion, due to the semi-insoluble nature of fatty acids

⁹ Short and medium-chain fatty acids can transport through the membranes freely, and therefore are not subject to this system. This also means that they are not subject to regulation via malonyl-CoA-dependent inhibition.




Figure 1: Schematic of the regulation of fatty acid oxidation. ACS indicates Acyl-CoA Synthetase, ACC indicates Acetyl-CoA Carboxylase.

¹⁰ Refer back to the lipid synthesis notes/lecture for more information

¹¹ Via activation by adrenaline or glucagon for example.

ited, Malonyl-CoA is reduced and CPTI is active. In this way, energy demand or adrenaline can promote fatty acid oxidation.

Lipid Oxidation in the Mitochondria

Once the Acyl-CoA is inside the mitochondria, oxidation occurs through a series of enzymatic reactions. These reactions proceed in a serial manner to extract one Acetyl-CoA at a time, removing two carbons each cycle. The cycles consist of the following:

Desaturation This generates a double bond at the Δ^2 position of the fatty acid¹². This step *generates* one FADH₂ molecule.

Isomerization This transfers the electrons from this new double bond to a new ketone group on the Δ^3 carbon. This step *generates* one NADH molecule.

Release The first two carbons are now released as Acetyl-CoA by an enzyme called a thiolase.

After one round, we are left with a fatty acid that is two carbons shorter, and have generated one Acetyl-CoA, one NADH and one FADH₂ that can be used as fuel. For a saturated fatty acid such as palmitate (C16:0) this means that this cycle occurs *seven* times¹³.

Alternative Fatty Acid Catabolism

Long chain, unsaturated fatty acids are catabolized in a very similar manner with one difference. The first desaturation step introduces a double bond, but for unsaturated fatty acids the double bond is already present. That means that during progressive oxidation, if a double bond is already present at the Δ^2 of the now shortened fatty acid the first step is skipped, and FADH₂ is not generated.

ODD-NUMBERED FATTY ACIDS after the last thiolase step, you end up with an Acetyl-CoA and a three carbon Propionyl-CoA¹⁴. Propionyl-CoA is then converted to Succinyl-CoA by a three step reaction that in net consumes one ATP and requires Vitamin B12. Succinyl-CoA is a component of the TCA cycle, and can be further oxidized generating one GTP and one NADH.

VERY LONG CHAIN FATTY ACIDS ARE FIRST OXIDIZED IN THE PEROXISOME. The Acyl-CoA Synthetase with activity towards very long chain fatty acids¹⁵ resides on the peroxisomes. Peroxisomes are membrane enclosed organelles that contain similar enzymes to those that perform β -oxidation in the mitochondria. The major difference is

¹² The Co-A group is attached at the carboxyl group, or the Δ^1 position.

¹³ Not eight, because the last reaction ends with two Acetyl-CoA molecules.

¹⁴ Several amino acids, including Isoleucine, Valine, Threonine and Methionine also produce Propionyl-CoA

¹⁵ Those with more than 22 carbons.

that while these lipids are catabolized, there is no electron transport chain in peroxisomes, so no ATP is produced. Instead electrons are transferred to peroxide, which is in turn converted to water and oxygen by the enzyme Catalase. This process terminates with short and medium chain fatty acids that are released as acyl-carnitines where they travel to mitochondria for final catabolism and some energy production.

Determining the Energy Content of a Fatty Acid

Based on the series of reactions above, or each two-carbon unit released, one Acetyl-CoA, one NADH and one FADH₂ are released. Added together this provides 14 ATP equivalents per two carbon units. Therefore to calculate the ATP in a saturated fatty acid like palmitate (C_{16:0}) we could do the following:

- Consume 2 ATP to activate the fatty acid (see the ACS reaction 1).
- Take the fatty acid length, and subtract 2 then divide by 2. This is the number of fatty acid oxidation cycles. For palmitate that is $\frac{16-2}{2} = 7$. For each cycle we generate 14 ATP units, for palmitate there is $14 \times 7 = 98$ ATP equivalents generated.
- The remaining Acetyl-CoA from the last two carbons left adds another 10 ATP equivalents
- The total ATP is then $98 + 10 - 2$ or 106 ATP equivalents.

For unsaturated fatty acids, you do not need to perform the desaturation step, so for each double bond there is one less FADH₂ generated than normal. If we had a fatty acid that was 16:1 it would generate a total of 104.5 ATP equivalents.

GENERALLY THIS PROCEDURE EXPLAINS TWO BIOENERGETIC PROPERTIES OF FATTY ACIDS. Longer chain fatty acids have more energy than shorter chain fatty acids, and saturated fatty acids have more energy than desaturated fatty acids. I have used the method above to calculate the ATP equivalents of several fatty acids in Table 1 to illustrate this point. Don't try to memorize these, but rather spend some time practicing how to calculate the ATP-equivalents¹⁶. After you have thought about how to calculate this, take a step back. This is a lot of energy storage. A triglyceride with three C_{16:0} molecules is catabolized to 318 ATP equivalents. This is close to *ten times* the energy storage of a fully oxidized glucose molecule (32 ATP).


Table 1: ATP equivalents of some fatty acids.

Lipid	Net ATP
Butyrate (C _{4:0})	22
Laurate (C _{12:0})	78
Palmitate (C _{16:0})	106
Oleate (C _{18:0})	120
Linoleate (C _{18:2})	117

¹⁶ For even more of a challenge, try to calculate the net energy production from an odd-chain fatty acid.

Ketolysis

Ketone bodies are produced when there is excessive acetyl-CoA, but insufficient TCA cycle intermediates in the liver. These ketone bodies¹⁷ can be converted back to Acetyl-CoA in non-hepatic tissues. This specificity is due to the absence of an enzyme called 3-oxoacid CoA-transferase¹⁸ in the liver thus not allowing the liver to utilize ketone bodies while allowing other tissues to utilize ketone bodies for energy. This enzyme catalyzes this reaction starting with Acetoacetate¹⁹:



As a result this generates a single Acetyl-CoA molecule after a thiolase reaction. This reaction is not cataplerotic because both Succinate and Succinyl-CoA are part of the TCA cycle. However bypassing the Succinyl-CoA Synthetase step means that one GTP is not generated during normal TCA cycle progress. Overall, an Acetyl-CoA that is released from the liver as β -hydroxybutyrate costs the liver 3.5 ATP molecules²⁰. In the recipient tissue, that β -hydroxybutyrate generates 6.5 ATP in net²¹. Overall, ketone usage means that the original Acetyl-CoA in the liver now only generates 3 ATP equivalents rather than the 10 ATP if it remained in the liver (6.5 ATP generated - 3.5 ATP used in liver to generate the β -hydroxybutyrate = 3 ATP in total). Table 2 summarizes the energy costs of the three major transport processes we have discussed this year. The *inefficiency* of these transport processes comes at a cost to performance, but can cause negative energy balance for weight loss. In terms of low-carbohydrate, high fat diets this may explain the slight increase in energy expenditure (100-200kcal/day) observed in relation to low fat, high carbohydrate diets [Hall et al., 2016, Ebbeling et al., 2018].

Regulation of Fatty Acid Oxidation

Short-term Regulation

As we described above, short term regulation is accomplished primarily through the ACC/Malonyl-CoA/CPTI system described in Figure 1. Two other key regulatory systems to consider in the short term are the flux of fatty acids, either via the regulation of lipolysis, or the regulation of Lipoprotein Lipase activity²². Another regulatory mechanism that might not seem as apparent, is that when there is sufficient non-lipid fuel sources (for example an excess of carbohydrates or amino acids), there will be more Malonyl-CoA and therefore less lipid oxidation. Finally, recall that the products of fatty

¹⁷ Primarily β -hydroxybutyrate then acetoacetate; acetone is a terminal end product and cannot be used for energy.

¹⁸ OXCT1, sometimes called succinyl-CoA-3-oxaloacid CoA transferase (or SCOT).

¹⁹ β -hydroxybutyrate can be interconverted to and from acetoacetate using up one NADH molecule for each acetoacetate generating, and using an NADH for the reverse reaction.

²⁰ 1 ATP for the ATP-Citrate Lyase step, 2.5 ATP for the NADH used to generate β -hydroxybutyrate

²¹ Generate 10 ATP from the TCA cycle oxidation of Acetyl-CoA, but using 1 NADH converting β -hydroxybutyrate to acetoacetate; and 1 GTP skipping the Succinyl-CoA Synthetase step. So $10 - 3.5 = 6.5$

Table 2: Energy costs of the macromolecule transport processes discussed this year. Refer to the gluconeogenesis notes for details about the Cori and Cahill cycles.

Process	Net ATP Loss
Ketone Body Transport	-7 ATP
Cori Cycle (Lactate)	-6 ATP
Cahill Cycle (Alanine)	-15 ATP

²² Discussed in the lipid transport lecture.

acid oxidation all require TCA cycle and electron transport chain activity. Since the most important driver of the electron transport chain is energy demand, the final oxidation of all that Acetyl-CoA, NADH and FADH₂ will only occur if there is some energy demand. Alternately, if there is insufficient TCA cycle intermediates²³ the NADH and FADH₂ generated from the partial oxidation of fatty acids might be used for fuel, but the Acetyl-CoA generated from the partial oxidation of fatty acids will undergo ketogenesis for transport to other tissues.

²³ For example due to cataplerosis in a liver undergoing substantial gluconeogenesis.

Transcriptional Adaptations for Fatty Acid Oxidation

IN TERMS OF ATHLETIC PERFORMANCE, increasing the ability to oxidize fatty acids is important for endurance athletes. Highly trained athletes tend to have more mitochondria, and higher expression of enzymes such as Lipoprotein Lipase, CPT₁, Acetyl-CoA Synthetase and enzymes of the TCA cycle. One major regulator of these adaptations are transcription factors called PPAR δ and PPAR α . In muscle tissues PPAR α and PPAR δ activate the transcription of several lipid transport and oxidation genes. In liver tissues, a similar transcription factor, PPAR α upregulates many of the same genes²⁴ along with genes involved in ketogenesis [Kersten et al., 2000, Badman et al., 2007]. Both of these transcription factors are induced by elevated intracellular fatty acid levels [Keller et al., 1993]. The specific natural ligands for the PPAR transcription factors have been difficult to unambiguously identify, but PUFA's seem to play a key role in activating these transcription factors [Forman et al., 1997]. This is potentially one mechanism by which PUFA's alleviate lipid accumulation.

²⁴ Activation of PPAR α is a promising target to alleviate hepatic steatosis and metabolic dysfunction-associated steatotic liver disease (MASLD), though they have not yet reached the clinic.

Interestingly, similar adaptations occur in the context of obesity, likely for the same reasons. The major difference, is that in the absence of energy demand, lipids are only partially oxidized or end up stored as lipids in cells. The buildup of stored, partially oxidized fatty acids is thought to play a role in the pathogenesis of insulin resistance, though the mechanisms are not clearly defined.

Reflection Questions

1. Calculate the net ATP yield from complete oxidation of one molecule of oleate (C₁₈:1 Δ^9) in muscle, accounting for the activation cost and the reduced FADH₂ generation due to the double bond. Then compare this to the net ATP a recipient muscle cell would obtain if the liver had instead exported that same Acetyl-

CoA as β -hydroxybutyrate. Use this comparison to evaluate the energetic trade-off of ketone body transport.

2. A patient with a primary carnitine deficiency cannot maintain adequate intracellular carnitine levels. Analyze how this would differentially affect oxidation of short-chain vs. long-chain fatty acids, predict the consequences during prolonged fasting, and explain why the heart would be particularly vulnerable in this condition.
3. A sedentary person and an endurance-trained athlete consume identical high-fat meals after an overnight fast. Evaluate how their capacity to oxidize dietary fatty acids differs at three levels: (a) hormonal control of lipolysis in adipose, (b) CPTI activity and malonyl-CoA regulation in muscle, and (c) transcriptional adaptations via PPAR α /PPAR δ . Predict the metabolic fate of dietary fatty acids in each person.

References

- Kevin J. Acheson, Yves Schutz, T. Bessard, K. Anantharaman, J. P. Flatt, and E. Jequier. Glycoprotein storage capacity and de novo lipogenesis during massive carbohydrate overfeeding in man. *American Journal of Clinical Nutrition*, 48(2):240–247, 1988. ISSN 00029165.
- Michael K. Badman, Pavlos Pissios, Adam R. Kennedy, George Koukos, Jeffrey S. Flier, and Eleftheria Maratos-Flier. Hepatic Fibroblast Growth Factor 21 Is Regulated by PPAR α and Is a Key Mediator of Hepatic Lipid Metabolism in Ketotic States. *Cell Metabolism*, 5(6):426–437, June 2007. ISSN 15504131. DOI: 10.1016/j.cmet.2007.05.002.
- Cara B. Ebbeling, Henry A Feldman, Gloria L Klein, Julia M W Wong, Lisa Bielak, Sarah K Steltz, Patricia K Luoto, Robert R Wolfe, William W Wong, and David S. Ludwig. Effects of a low carbohydrate diet on energy expenditure during weight loss maintenance: Randomized trial. *Bmj*, 363:k4583, 2018. ISSN 0959-8138. DOI: 10.1136/BMJ.K4583.
- J. M. Ellis, S. M. Mentock, M. a. DePetrillo, T. R. Koves, S. Sen, S. M. Watkins, D. M. Muoio, Gary W. Cline, H. Taegtmeier, Gerald I. Shulman, M. S. Willis, and Rosalind A. Coleman. Mouse Cardiac Acyl Coenzyme A Synthetase 1 Deficiency Impairs Fatty Acid Oxidation and Induces Cardiac Hypertrophy. *Molecular and Cellular Biology*, 31(6):1252–1262, January 2011. ISSN 0270-7306. DOI: 10.1128/MCB.01085-10.

- B M Forman, J Chen, and Ronald M. Evans. Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta. *Proceedings of the National Academy of Sciences of the United States of America*, 94(9):4312–7, April 1997. ISSN 0027-8424.
- Trisha J. Grevenkoed, Eric L Klett, and Rosalind A. Coleman. Acyl-CoA Metabolism and Partitioning. *Annual review of nutrition*, 34:1–30, 2014. ISSN 1545-4312. DOI: 10.1146/annurev-nutr-071813-105541.
- Kevin D. Hall, Kong Y Chen, Juen Guo, Yan Y Lam, Rudolph L. Leibel, Laurel ES Mayer, Marc L. Reitman, Michael Rosenbaum, Steven R Smith, B Timothy Walsh, and Eric Ravussin. Energy expenditure and body composition changes after an isocaloric ketogenic diet in overweight and obese men. *The American Journal of Clinical Nutrition*, 104(2):324–333, August 2016. ISSN 0002-9165. DOI: 10.3945/ajcn.116.133561.
- H Keller, C Dreyer, J Medin, A Mahfoudi, K Ozato, and W Wahli. Fatty acids and retinoids control lipid metabolism through activation of peroxisome proliferator-activated receptor-retinoid X receptor heterodimers. *Proceedings of the National Academy of Sciences of the United States of America*, 90(6):2160–4, 1993. ISSN 0027-8424. DOI: 10.1073/pnas.90.6.2160.
- Sander Kersten, Béatrice Desvergne, and Walter Wahli. Roles of PPARs in health and disease. *Nature*, 405(6785):421–424, 2000. ISSN 00280836. DOI: 10.1038/35013000.
- Kwangwon Lee, Janos Kerner, and Charles L. Hoppel. Mitochondrial carnitine palmitoyltransferase 1a (CPT1a) is part of an outer membrane fatty acid transfer complex. *Journal of Biological Chemistry*, 286(29):25655–25662, 2011. ISSN 00219258. DOI: 10.1074/jbc.M111.228692.
- R. M. McDevitt, S. J. Bott, M. Harding, W. A. Coward, L. J. Bluck, and A. M. Prentice. De novo lipogenesis during controlled overfeeding with sucrose or glucose in lean and obese women. *American Journal of Clinical Nutrition*, 74(6):737–746, 2001. ISSN 00029165. DOI: 10.1056/nejm-jw.FW109926.
- J D McGarry, G P Mannaerts, and D W Foster. A possible role for malonyl-CoA in the regulation of hepatic fatty acid oxidation and ketogenesis. *The Journal of clinical investigation*, 60(1):265–70, July 1977. ISSN 0021-9738. DOI: 10.1172/JCI108764.

J R Neely and H E Morgan. Relationship Between Carbohydrate and Lipid Metabolism and the Energy Balance of Heart Muscle. *Annual Review of Physiology*, 36(1):413–459, March 1974. ISSN 0066-4278. DOI: 10.1146/annurev.ph.36.030174.002213.

John L Tymoczko, Jeremy M Berg, and Lubert Stryer. *Biochemistry: A Short Course*. W.H. Freeman and Co, New York, NY, 2015.